

Name: Jiangxi Wellanimal Gene Technology Co., Ltd. Address: 5F, Building 2, Ganzhou National High-level Talents Science and Innovation Park(Phase II), 1 Wudang Mountain Road, Zhanggong District High-tech zone, Ganzhou City, Jiangxi Province.

Telephone: 400-6723-898 E-mail: biolight_info@163.com Website: http://www.wellanimal.com

Anti-AMACR Antibody, Mouse Monoclonal, A1T

2-methylacyl-CoA racemase

Catalog: MA01524HuM10-A1T Size:100µL

Basic Info Host Species Reactivity

Mouse Human/Mouse/Monkey

Conjugate Clonality, Isotype

None /

Concentration Immunogen

1mg/mL A synthetic peptide of human AMACR

Physical State Purification

Liquid Immunogen affinity purified

Clone No Applications

A1T WB/IHC

Property Form & Buffer: Supplied in PBS, 50% glycerol, PH7.4.

Specificity / Sensitivity: MA01524HuM10-A1T recognizes endogenous levels

of human AMACR.

Applications The application notes include recommended starting dilutions;

optimal dilutions/concentrations should be determined by the end user.

WB: $0.5 \sim 5 \mu g/mL$ IF: $5 \sim 20 \mu g/mL$ IHC: $5 \sim 20 \mu g/mL$ ICC: $5 \sim 20 \mu g/mL$

Usage and

Store at $4^{\circ}C$ for frequent use.

Storage

Aliquot and store at -20℃ for 12 months.

Avoid repeated freezing/thawing and violent shaking.

Please centrifuge it, before using.

QC Data




Figure 1. Application in WB

Western blot analysis of extracts of various cell lines and tissues, using AMACR antibody MAO1524HuM10-A1T at $2\mu g/mL$ Secondary antibody: HRP Mouse Anti-Rabbit IgG (H+L) at 1: 5000 dilution. Lysates/proteins: $25\mu g$ per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 2s.



QC Data




Figure 2. Application in IHC

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of monkey liver, monkey kidney sections labelling AMACR with purified MA01524HuM10-A1T at $30\mu g/mL$ dilution. Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using citrate buffer (PH6.0). Tissue was counterstained with Hematoxylin. Mouse specific IHC polymer detection kit HRP/DAB secondary antibody was used at 1: 2000 dilution. PBS instead of the primary antibody was used as the negative control.

For Research Use Only!