

Anti-TP53 Antibody, Rabbit Polyclonal

Synonyms: TP53, LFS1, TRP53, Li-Fraumeni Syndrome, Cellular tumor antigen p53, Antigen NY-C0-13, Phosphoprotein p53, Tumor suppressor p53

Basic Info

Catalog

PA00116HuA10

Host

Rabbit

Conjugate

None

Size

100 μ L

Concentration

1.0 mg/mL

Physical State

Liquid

Species Reactivity

Predict reacts with: Human/Mouse/Rat

Immunogen

Recombinant human TP53 protein, fragment

Met1~Asp393;

UniprotKB: P04637

Purification

Antigen Affinity Chromatography

Applications

WB/IHC

Property

Form & Buffer: Supplied in PBS, 50% glycerol, PH7.4.

Specificity / Sensitivity: Anti-TP53 Antibody, Rabbit Polyclonal recognizes endogenous levels of total TP53 protein.

Usage and Storage

Shipped at 4°C.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Avoid repeated freezing/thawing and violent shaking.

Please centrifuge it, before using.

Applications

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB: 0.5~5 μ g/mL

IF: 5~20 μ g/mL

IHC: 5~20 μ g/mL

ICC: 5~20 μ g/mL

QC Data




Figure 1. Application in WB

Western blot analysis of extracts of MCF7 cell line, using TP53 antibody (PA00116HuA10) at 1 μ g/mL. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1 s.

QC Data




Figure 2. Application in IHC

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon sections labelling TP53 with purified PA00116HuA10 at 10 μ g/mL. Heat mediated antigen retrieval was performed using citrate buffer (pH 6.0). Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB secondary antibody was used at 1/4000 dilution. PBS instead of the primary antibody was used as the negative control.

For Research Use Only!